arXiv:1602.06616v1 [math.DG] 22 Feb 2016

SOME FUNCTIONALS ON COMPACT MANIFOLDS WITH BOUNDARY

PENGZI MIAO[1] AND LUEN-FAI TAM[2]

Abstract. We analyze critical points of two functionals of Riemannian metrics on compact manifolds with boundary. These functionals are motivated by formulae of the mass functionals of asymptotically flat and asymptotically hyperbolic manifolds

1. introduction

In this work we discuss critical points of two functionals defined on the space of Riemannian metrics on a compact manifold with boundary. Let Ω be an n-dimensional, connected, compact manifold with smooth boundary Σ which itself is not necessarily connected. Given a Riemannian metric g on Ω, let

$$G_\lambda^g := \mathrm{Ric}(g) - \frac{1}{2}\left[\mathcal{S}_g - \lambda(n-1)(n-2)\right] g, \tag{1.1}$$

where $\mathrm{Ric}(g)$, $\mathcal{S}_g$ are the Ricci curvature, the scalar curvature of g, respectively, and $\lambda \in \{0, 1, -1\}$. Let X be a vector field on Ω. Define

$$\mathcal{F}_X(g) := \int_\Sigma G_\lambda^g(X, \nu_g) d\gamma_g, \tag{1.2}$$

where ν_g is the outward unit normal to Σ in (Ω, g) and $d\gamma_g$ denotes the volume element on Σ induced by g.

We will be interested in $\mathcal{F}_X(\cdot)$ constrained to the space of constant scalar curvature metrics on Ω. To be precise, we assume that g_0 is a given, smooth metric on Ω such that $\mathcal{S}_{g_0} = \lambda n(n-1)$. If $\lambda > 0$, we assume that the first Dirichlet eigenvalue of $(n-1)\Delta_{g_0} + \mathcal{S}_{g_0}$ is positive.

Date: February, 2016.
2010 *Mathematics Subject Classification.* Primary 53C20; Secondary 83C99.
Key words and phrases. Scalar curvature, static metrics.
[1]Research partially supported by Simons Foundation Collaboration Grant for Mathematicians #281105.
[2]Research partially supported by Hong Kong RGC General Research Fund #CUHK 14305114.

Let γ be the metric on Σ induced from g_0. Define

$$\mathcal{M}^\lambda := \{g \text{ is a smooth Riemannian metric on } \Omega \text{ with } \mathcal{S}_g = \lambda n(n-1)\},$$
$$\mathcal{M}^\lambda_\gamma := \{g \in \mathcal{M}^\lambda \text{ such that } g|_{T(\Sigma)} = g_0|_{T(\Sigma)}\},$$
$$\mathcal{M}^\lambda_0 := \{g \in \mathcal{M}^\lambda \text{ such that } g = g_0 \text{ everywhere at } \Sigma\}.$$

Here, in the definition of $\mathcal{M}^\lambda_\gamma$, $T(\Sigma)$ denotes the tangent bundle of Σ. Evidently, $g_0 \in \mathcal{M}^\lambda$ and $\mathcal{M}^\lambda_0 \subset \mathcal{M}^\lambda_\gamma \subset \mathcal{M}^\lambda$.

To state our result, we will assume X is a *conformal Killing* vector field on (Ω, g_0), see **(a2)** in section 2. Under this assumption, it is known (cf. [6]) that

$$g_0 \text{ is an Einstein metric} \Rightarrow \mathrm{div}_{g_0} X \in \mathrm{Ker}(D\mathcal{S}^*_{g_0}),$$

where "div_{g_0}" is the divergence with respect to g_0 and $\mathrm{Ker}(D\mathcal{S}^*_{g_0})$ denotes the kernel of the operator $D\mathcal{S}^*_{g_0}(\cdot)$, which is the formal L^2-adjoint of the linearization of the scalar curvature map $\mathcal{S}$ at g_0, i.e.

$$D\mathcal{S}^*_{g_0}(f) = -(\Delta_{g_0} f)g_0 + \nabla^2_{g_0} f - f\mathrm{Ric}(g_0)$$

for any function f. In a relativistic context, a nontrivial solution f to $D\mathcal{S}^*_{g_0}(f) = 0$ is often referred as a static potential. Motivated by the work of Corvino [3] on localized scalar curvature deformation, we seek a variational characterization of the condition "$\mathrm{div}_{g_0} X \in \mathrm{Ker}(D\mathcal{S}^*_{g_0})$", which is part of the following theorem.

Theorem 1.1. *Let (Ω, g_0), $\mathcal{M}^\lambda$, $\mathcal{M}^\lambda_\gamma$ and $\mathcal{M}^\lambda_0$ be described as above. Suppose X is a conformal Killing vector field on (Ω, g_0). Let $\{g(t)\}_{|t|<\epsilon}$ be a smooth path of metrics on Ω such that $g(0) = g_0$. Then*

(i) $\dfrac{d}{dt}\mathcal{F}_X(g(t))|_{t=0} = 0$ *for all path $\{g(t)\} \subset \mathcal{M}^\lambda_0$ if and only if $div_{g_0} X \in \mathrm{Ker}(D\mathcal{S}^*_{g_0}(\cdot))$.*

(ii) $\dfrac{d}{dt}\mathcal{F}_X(g(t))|_{t=0} = 0$ *for all path $\{g(t)\} \subset \mathcal{M}^\lambda_\gamma$ if and only if $div_{g_0} X \in \mathrm{Ker}(D\mathcal{S}^*_{g_0}(\cdot))$ and ν_{g_0} is an eigenvector of $\langle X, \nu_{g_0}\rangle G^{g_0}$ with zero eigenvalue at every point on Σ.*

(iii) *If g_0 is Einstein, then $\dfrac{d}{dt}\mathcal{F}_X(g(t))|_{t=0} = 0$ for all path $\{g(t)\} \subset \mathcal{M}^\lambda$. The converse is true if there is a function u such that $div_{g_0} X - X(u)$ has a fixed sign on a dense subset in Ω, which is satisfied if $div_{g_0} X$ does not vanish on Ω.*

The use of the functional $\mathcal{F}_X(\cdot)$ in Theorem 1.1 is largely motivated by corresponding expressions of the mass functionals of asymptotically

flat and asymptotically hyperbolic manifolds. On an asymptotically flat manifold (M^n, g), the ADM mass [1] $\mathfrak{m}_{ADM}$ can be computed by

$$\mathfrak{m}_{ADM} = c_n \lim_{r\to\infty} \int_{S_r} G_0^g(X, \nu_g) d\gamma, \tag{1.3}$$

where $c_n < 0$ is a dimensional constant, S_r denotes the coordinate sphere of radius r in the asymptotically flat end of (M, g), and

$$X = \sum_{i=1}^{n} x^i \frac{\partial}{\partial x^i}$$

is the conformal Killing vector field on the Euclidean space $\mathbb{R}^n$. For the origin and the proof of (1.3), we refer readers to [2, 4, 7, 12, 6, 14]. On an asymptotically hyperbolic manifold, formulae analogous to (1.3), which compute the mass functionals introduced in [5], can be found in [6].

The second functional that we are interested in is also related to (1.3). Indeed, it is not hard to see that one can replace $X = \sum_i x^i \frac{\partial}{\partial x^i}$ in (1.3) by $X = r\nu_g$ (cf. [13]) to obtain

$$\begin{aligned} \mathfrak{m}_{ADM} &= c_n \lim_{r\to\infty} \int_{S_r} r \left[\mathrm{Ric}(g) - \frac{1}{2}\mathcal{S}_g \right] (\nu_g, \nu_g) d\gamma \\ &= c_n \lim_{r\to\infty} \int_{S_r} r(H_2 - H_2^0) d\gamma, \end{aligned} \tag{1.4}$$

where one has also used the Gauss equation and H_2, H_2^0 denote the second order mean curvature of S_r in (M, g), $\mathbb{R}^n$, respectively. (Here one assumes that S_r can be isometrically embedded in $\mathbb{R}^n$ which is always satisfied when $n = 3$). We recall that the second order mean curvature H_2 of a hyersurface S in a Riemannian manifold (M, g) is given by

$$H_2 := \sum_{1\le\alpha<\beta\le n-1} \kappa_\alpha \kappa_\beta = \frac{1}{2}\left(H^2 - |A|^2\right), \tag{1.5}$$

where $\{\kappa_\alpha\}_{1\le\alpha\le n-1}$, A and H are the principal curvatures, the second fundamental form and the (first order) mean curvature of S in (M, g), respectively. For a generalization of (1.4), we refer readers to Theorem 1.2 in [13].

Motivated by (1.4), given a nontrivial function ϕ on $\Sigma = \partial\Omega$, we define

$$\mathcal{E}_\phi(g) := \int_\Sigma \phi\, H_2(g) d\gamma, \tag{1.6}$$

where $H_2(g)$ is the second order mean curvature of Σ in (Ω, g) with resect to ν_g. Regarding $\mathcal{E}_\phi(\cdot)$, we have

Theorem 1.2. *Let (Ω, g_0) and $\mathcal{M}_\gamma^\lambda$ be described as above. Let H_{g_0} be the mean curvature of Σ in (Ω, g_0). Let $\{g(t)\}_{|t|<\epsilon}$ denote a smooth path of metrics on Ω such that $g(0) = g_0$. Then*

$$\frac{d}{dt}\mathcal{E}_\phi(g(t))|_{t=0} = 0 \quad \textit{for all } \{g(t)\} \subset \mathcal{M}_\gamma^\lambda$$

if and only if the following conditions hold

(i) *$\phi\, H_{g_0}$ is the boundary value of some $N \in \mathrm{Ker}(D\mathcal{S}_{g_0}^*)$.*
(ii) *Σ is umbilical at every point $p \in \Sigma$ where $\phi(p) \neq 0$.*

We give some remarks about Theorem 1.2.

Remark 1.1. Besides (1.4), our motivation to study $\mathcal{E}_\phi(\cdot)$ on $\mathcal{M}_\gamma^\lambda$ also comes from a corresponding result on a weighted total mean curvature functional in [10]. More precisely, if we let

$$\mathcal{E}_\phi^{(1)}(g) = \int_\Sigma \phi H(g) d\gamma_g,$$

where $H(g)$ is the usual mean curvature of Σ in (Ω, g), it was proved in [10] that g_0 is a critical point of $\mathcal{E}_\phi^{(1)}(\cdot)$ in $\mathcal{M}_\gamma^\lambda$ if and only if ϕ is the boundary value of some element $N \in \mathrm{Ker}(D\mathcal{S}_{g_0}^*)$.

Remark 1.2. If (Ω, g_0) has a conformal Killing vector field X such that X does not vanish along Σ and is orthogonal to Σ, then every point in Σ is umbilical (see Proposition 4.2 in Section 4).

This paper is organized as follows. In Section 2, we discuss properties of manifolds of constant scalar curvature admitting a conformal Killing vector field. In Section 3, we prove Theorem 1.1. In Section 4, we prove Theorem 1.2.

2. Manifolds with a conformal Killing vector field

Let Ω, Σ be given as in the introduction. Let g_0 be a Riemannian metric on Ω satisfying

(a1): g_0 has constant scalar curvature $\lambda n(n-1)$, with $\lambda = 0, 1$ or -1, where n is the dimension of Ω;

(a2): (Ω, g_0) has a conformal Killing vector field X, i.e.

$$\eta_{i|j} + \eta_{j|i} = \frac{2}{n}(\mathrm{div}_{g_0} X)(g_0)_{ij}$$

where η is the 1-form dual to X with respect to g_0 and " $_|$ " denotes the covariant differentiation on (Ω, g_0).

Lemma 2.1. *Suppose (Ω, g_0) satisfies conditions* **(a1)**, **(a2)**. *Then*

(i) *$tr_{g_0}(D\mathcal{S}^*_{g_0}(div_{g_0}X)) = 0$, where "$tr_{g_0}(\cdot)$" denotes the trace with respect to g_0.*

(ii) *$D\mathcal{S}^*_{g_0}(div_{g_0}X) = -\frac{n}{n-2}\left[L_X G^{g_0}_\lambda + \frac{n-2}{n}(div_{g_0}X)G^{g_0}_\lambda\right]$, where "$L_X$" denotes the Lie derivative along X.*

Proof. For simplicity, we let $f = \mathrm{div}_{g_0}X$ and denote $G^{g_0}_\lambda$ by G. For any smooth symmetric $(0,2)$ tensor h, computing in a local orthonormal frame $\{e_i\}_{1\le i\le n}$, we have

$$\begin{aligned}
&\langle L_X G, h\rangle - \langle G, L_X h\rangle \\
&= \langle \nabla_X G, h\rangle - \langle G, \nabla_X h\rangle + \left(G_{kj}\eta_{k|i} + G_{ik}\eta_{k|j}\right)h_{ij} - \left(h_{kj}\eta_{k|i} + h_{ik}\eta_{k|j}\right)G_{ij} \\
&= 2\langle \nabla_X G, h\rangle - X\langle G, h\rangle + h_{ij}\left(G_{kj}\eta_{k|i} + G_{ik}\eta_{k|j}\right) - h_{ij}\eta_{i|k}G_{kj} - h_{ij}\eta_{j|k}G_{ik} \\
&= 2\langle \nabla_X G, h\rangle - X\langle G, h\rangle + h_{ij}\left[2\left(G_{kj}\eta_{k|i} + G_{ik}\eta_{k|j}\right) - \frac{4}{n}fG_{ij}\right] \\
&= 2\langle L_X G, h\rangle - \frac{4}{n}f\langle G, h\rangle - X\langle G, h\rangle,
\end{aligned}$$

where summation is taken over any pair of repeated indices and **(a2)** is applied. Therefore,

$$\langle L_X G, h\rangle = -\langle G, L_X h\rangle + \frac{4}{n}f\langle G, h\rangle + X\langle G, h\rangle. \tag{2.1}$$

Now let $h = g_0$. By (2.1), **(a2)** and the fact $\mathcal{S}_{g_0} = \lambda n(n-1)$,

$$\langle G, g_0\rangle = 0 \quad \text{and} \quad \mathrm{tr}_{g_0}(L_X G) = 0. \tag{2.2}$$

To proceed, we adopt an argument of Herzlich [6]. Let $\{\phi_t\}_{|t|<\epsilon}$ be the 1-parameter family of local diffeomorphisms generated by X. Let $g_t = \phi_t^*(g_0)$, then $g_t = e^{2u_t}g_0$ for some smooth function $u(\cdot, t) = u_t(\cdot)$ with $u_0 = 0$. Let $v = \frac{\partial u}{\partial t}|_{t=0}$. By **(a2)** and the fact $L_X g_0 = \frac{d}{dt}g_t|_{t=0}$,

$$f = nv. \tag{2.3}$$

On the other hand,

$$\begin{aligned}
\mathrm{Ric}(g_t) = \mathrm{Ric}(g_0) &- (n-2)(\nabla^2_{g_0}u_t - du_t \otimes du_t) \\
&- \left(\Delta_{g_0}u_t + (n-2)|du_t|^2_{g_0}\right)g_0.
\end{aligned} \tag{2.4}$$

Taking the t-derivative and letting $t = 0$, we have

$$L_X\mathrm{Ric}(g_0) = -(n-2)\nabla^2_{g_0}v - (\Delta_{g_0}v)g_0. \tag{2.5}$$

Hence, by (2.3) and (2.5),

$$\begin{aligned} & L_X G + \frac{n-2}{n} f G \\ = & -(n-2) D\mathcal{S}^*_{g_0}(v) - (n-1)\left(\Delta_{g_0} v + \lambda n v\right) g_0 \\ = & -\frac{n-2}{n} D\mathcal{S}^*_{g_0}(f) + \frac{1}{n}\mathrm{tr}_{g_0}\left(D\mathcal{S}^*_{g_0}(f)\right) g_0. \end{aligned} \tag{2.6}$$

Taking trace of (2.6) and using (2.2), we conclude that (i) and (ii) of the lemma hold. □

If g_0 is Einstein, then $G^{g_0}_\lambda = 0$, which implies $D\mathcal{S}^*_{g_0}(\mathrm{div}_{g_0} X) = 0$ by Lemma 2.1 (ii). (This was the content of [6, Lemma 2.2].) Next we show that the reverse is true under certain boundary and interior conditions.

Proposition 2.1. *Let (Ω, g_0) satisfying conditions* **(a1), (a2)**. *Suppose*

(i) $D\mathcal{S}^*_{g_0}(div_{g_0} X) = 0$,
(ii) $\langle X, \nu_{g_0}\rangle |G^{g_0}|^2 \geq 0$ *along* Σ, *and*
(iii) *there exists a smooth function u on Ω such that $div_{g_0} X > X(u)$ on a dense subset of Ω.*

Then g_0 is Einstein in Ω.

Proof. Denote $G^{g_0}_\lambda$ by G. By (i) and Lemma 2.1,

$$L_X G + \frac{n-2}{n}(\mathrm{div}_{g_0} X) G = 0.$$

Therefore,

$$\begin{aligned} \frac{1}{2} X(|G|^2) &= \langle \nabla_X G, G\rangle \\ &= \langle L_X G, G\rangle - G_{kj}\eta_{k|i} G_{ij} - G_{ki}\eta_{k|j} G_{ij} \\ &= -\frac{n-2}{n}(\mathrm{div}_{g_0} X)|G|^2 - \frac{2}{n}(\mathrm{div}_{g_0} X)|G|^2 \\ &= -(\mathrm{div}_{g_0} X)|G|^2. \end{aligned} \tag{2.7}$$

Multiplying (2.7) by a function ϕ and integrating on Ω, we have

$$\begin{aligned} & -2\int_\Omega \phi(\mathrm{div}_{g_0} X)|G|^2 dV_{g_0} \\ = & \int_\Omega \left[X(\phi|G|^2) - X(\phi)|G|^2\right] dV_{g_0} \\ = & \int_\Omega \left[\mathrm{div}_{g_0}(\phi|G|^2 X) - \phi \mathrm{div}_{g_0}(X)|G|^2 - X(\phi)|G|^2\right] dV_{g_0}, \end{aligned}$$

where dV_{g_0} denotes the volume element of g_0. Hence,

$$\int_\Omega \left[X(\phi) - \phi \mathrm{div}_{g_0}(X)\right] |G|^2 dV_{g_0} = \int_\Sigma \phi |G|^2 \langle X, \nu_{g_0}\rangle d\gamma_{g_0}.$$

Letting $\phi = e^u$, by (ii) we have

$$\int_\Omega e^u \left[X(u) - \mathrm{div}_{g_0}(X)\right] |G|^2 dV_{g_0} \geq 0.$$

By (iii), we conclude that $G \equiv 0$ and g_0 is Einstein. □

Remark 2.1. Proposition 2.1 implies that, if $\mathrm{div}_{g_0} X \in \mathrm{Ker}(D\mathcal{S}^*_{g_0})$ with $\langle X, \nu_{g_0}\rangle \geq 0$ along Σ and $\mathrm{div}_{g_0} X > 0$ in Ω, then g_0 is Einstein.

3. Critical points of $\mathcal{F}_X(\cdot)$

Let g_0 be a metric on Ω satisfying conditions **(a1), (a2)**. In this section, we consider the functional $\mathcal{F}_X(\cdot)$ near g_0.

Lemma 3.1. *Let $\{g(t)\}_{|t|<\epsilon}$ be a smooth family of metrics on Ω with $g(0) = g_0$ and $g'(0) = h$.*

(i) *The derivative of $\mathcal{F}_X(g(t))$ at $t = 0$ is given by*

$$\begin{aligned}&\frac{d}{dt}\mathcal{F}_X(g(t))|_{t=0}\\ &= \frac{n-2}{2n}\int_\Omega \left[\langle D\mathcal{S}^*_{g_0}(div_{g_0}X), h\rangle_{g_0} - (div_{g_0}X) D\mathcal{S}_{g_0}(h)\right] dV_{g_0}\\ &\quad + \frac{1}{2}\int_\Sigma \langle G^{g_0}_\lambda, h\rangle_{g_0} \langle X, \nu_{g_0}\rangle_{g_0} d\gamma_{g_0},\end{aligned}$$

where $D\mathcal{S}_{g_0}(\cdot)$ is the linearization of the scalar curvature at g_0.

(ii) *If $h|_{T(\Sigma)} = 0$, then*

$$\begin{aligned}&\frac{d}{dt}\mathcal{F}_X(g(t))|_{t=0}\\ &= \int_\Sigma \left[\frac{n-2}{n}(div_{g_0}X) DH_{g_0}(h) + \frac{1}{2}\langle G^{g_0}_\lambda, h\rangle_{g_0}\langle X, \nu_{g_0}\rangle_{g_0}\right] d\gamma_{g_0},\end{aligned}$$

where $DH_{g_0}(\cdot)$ is the linearization of the mean curvature H of Σ at g_0.

Proof. Let $\eta(t)$ be the 1-form dual to X with respect to $g(t)$. Let " ; " denote the covariant derivative on $(\Omega, g(t))$. Let $dV_{g(t)}$ be the volume element of $g(t)$. Denote $G^{g(t)}_\lambda$ by G^t. By the contracted Bianchi identity, $\mathrm{div}_{g(t)} G^t = 0$. Hence, $\mathcal{F}_X(g(t))$ can be rewritten as

$$\mathcal{F}_X(g(t)) = \int_\Omega \langle G^t, \beta^t\rangle_{g(t)} dV_{g(t)}, \tag{3.1}$$

where $\beta^t_{ij} = \frac{1}{2}(\eta_{i;j} + \eta_{j;i})$.

At $t = 0$, **(a1)** and **(a2)** imply

$$\langle G^0, \beta^0 \rangle_{g_0} = 0 \tag{3.2}$$

and

$$\frac{d}{dt}\langle G^t, \beta^t \rangle_{g(t)} = \frac{1}{n} f \left[-2\langle G^0, h\rangle_{g_0} + \mathrm{tr}_{g_0}(\frac{d}{dt}G^t) \right] + \langle G^0, \frac{d}{dt}\beta^t \rangle_{g_0}. \tag{3.3}$$

where $f = \mathrm{div}_{g_0} X$. Using the fact

$$\mathrm{tr}_{g(t)} G^t = \frac{n-2}{2}\left[-\mathcal{S}_{g(t)} + \lambda(n-1)n \right] \text{ and } \beta^t = \frac{1}{2} L_X g(t),$$

one has

$$\mathrm{tr}_{g_0}(\frac{d}{dt}G^t)|_{t=0} = -\frac{n-2}{2} D\mathcal{S}_{g_0}(h) + \langle G^0, h\rangle_{g_0} \tag{3.4}$$

and

$$\langle G^0, \frac{d}{dt}\beta^t \rangle_{g_0}|_{t=0} = \frac{1}{2}\langle G^0, L_X h\rangle_{g_0}. \tag{3.5}$$

Therefore, by (3.1) – (3.5) and (2.1), we have

$$\begin{aligned} &\frac{d}{dt}\mathcal{F}_X(g(t))|_{t=0} \\ =\ & \int_\Omega \left[-\frac{n-2}{2n} f D\mathcal{S}_{g_0}(h) + \frac{1}{2}\langle G^0, L_X h - \frac{2}{n} f h\rangle_{g_0} \right] dV_{g_0} \\ =\ & \int_\Omega \Big\{ -\frac{n-2}{2n} f D\mathcal{S}_{g_0}(h) + \frac{1}{2}\langle -L_X G^0 + \frac{2-n}{n} f G^0, h\rangle_{g_0} \\ & + \frac{1}{2}\mathrm{div}_{g_0}\left[\langle G^0, h\rangle_{g_0} X \right] \Big\} \end{aligned} \tag{3.6}$$

(i) now follows from (3.6) and Lemma 2.1(ii). When $h|_{T(\Sigma)} = 0$, one easily checks that the following formula

$$\int_\Omega w D\mathcal{S}_{g_0}(h) dV_{g_0} = \int_\Omega \langle D\mathcal{S}^*_{g_0}(w), h\rangle_{g_0} dV_{g_0} - 2\int_\Sigma w D H_{g_0}(h) d\gamma_{g_0} \tag{3.7}$$

holds for any function w on Ω (cf. (34) in [11] and (2.3) in [10]). Hence (ii) is true. □

Proof of Theorem 1.1. (i) For a variation $\{g(t)\} \subset \mathcal{M}^\lambda_0$, one has $h = 0$ at Σ and $D\mathcal{S}_{g_0}(h) = 0$ on Ω. Hence, by Lemma 3.1(i),

$$\frac{d}{dt}\mathcal{F}_X(g(t))|_{t=0} = \frac{n-2}{2n}\int_\Omega \langle D\mathcal{S}^*_{g_0}(\mathrm{div}_{g_0} X), h\rangle_{g_0} dV_{g_0}. \tag{3.8}$$

Therefore, if $\mathrm{div}_{g_0} X \in \mathrm{Ker}(D\mathcal{S}^*_{g_0})$, then $\frac{d}{dt}\mathcal{F}_X(g(t))|_{t=0} = 0$.

To see the converse is true, let $\hat{h}$ be an arbitrary symmetric $(0,2)$ tensor on Ω. Since the first Dirichlet eigenvalue of $(n-1)\Delta_{g_0} + \mathcal{S}_{g_0}$ is assumed to be positive, by the proof of [11, Proposition 1], one can find a positive function $u(x,t) = u_t(x)$, with $x \in \Omega$ and $|t| < \delta$ for some small δ, such that $u_0 = 1$ on Ω, $g(t) := u_t^{\frac{4}{n-2}}(g_0 + t\hat{h})$ has the same constant scalar curvature as g_0 and $u_t = 1$ at Σ for all t. Such a variation $\{g(t)\}$ satisfies

$$h := g'(0) = \frac{4}{n-2}\frac{\partial u}{\partial t}|_{t=0} g_0 + \hat{h} \tag{3.9}$$

with $h = \hat{h}$ at Σ. To proceed, we can choose $\hat{h}$ to have compact support in Ω for instance. Then, by (3.8), (3.9) and Lemma 2.1 (i),

$$0 = \frac{n-2}{2n}\int_\Omega \langle D\mathcal{S}^*_{g_0}(\mathrm{div}_{g_0} X), \hat{h}\rangle_{g_0} dV_{g_0},$$

which readily shows $D\mathcal{S}^*_{g_0}(\mathrm{div}_{g_0} X) = 0$.

(ii) Denote $G^{g_0}_\lambda$ by G^0 for simplicity. If $\mathrm{div}_{g_0} X \in \mathrm{Ker}(D\mathcal{S}^*_{g_0}(\cdot))$ and ν_{g_0} is an eigenvector of $\langle X, \nu_{g_0}\rangle G^0$ with zero eigenvalue everywhere at Σ, then $\frac{d}{dt}\mathcal{F}_X(g(t))|_{t=0} = 0$ by Lemma 3.1(i) and the fact $h|_{T(\Sigma)} = 0$.

Conversely, suppose $\frac{d}{dt}\mathcal{F}_X(g(t))|_{t=0} = 0$ for all $\{g(t)\} \subset \mathcal{M}^\lambda_\gamma$, then $\mathrm{div}_{g_0} X \in \mathrm{Ker}(D\mathcal{S}^*_{g_0})$ by (i). Now let $\hat{h}$ be any symmetric $(0,2)$ tensor with $\hat{h}|_{T(\Sigma)} = 0$. By the same construction of $\{g(t)\}$ used in (i), we have

$$0 = \frac{1}{2}\int_\Sigma \langle G^0, \hat{h}\rangle_{g_0} \langle X, \nu_{g_0}\rangle_{g_0} d\gamma_{g_0}. \tag{3.10}$$

Let $Q = \langle X, \nu_{g_0}\rangle G^0$ and choose $\hat{h}$ such that

$$\hat{h}(u,v) = Q(u,v) - Q(u^T, v^T) \ \text{ at } \Sigma,$$

where u, v are tangent vectors to Ω at Σ and u^T, v^T denote the project of u, v to the tangent space to Σ, respectively. With this choice of $\hat{h}$, (3.10) implies $Q(\nu_{g_0}, u) = 0$ for any u tangent to Ω at Σ. Therefore, ν_{g_0} is an eigenvector of Q with zero eigenvalue.

(iii) If g_0 is Einstein, then $G^0 = 0$ and $\mathrm{div}_{g_0} X \in \mathrm{Ker}(D\mathcal{S}^*_{g_0})$. Hence $\frac{d}{dt}\mathcal{F}_X(g(t))|_{t=0} = 0$ for all variation $\{g(t)\} \subset \mathcal{M}^\lambda$ by Lemma 3.1(i). Next, suppose $\frac{d}{dt}\mathcal{F}_X(g(t))|_{t=0} = 0$ for all such $\{g(t)\}$. Then $\mathrm{div}_{g_0} X \in \mathrm{Ker}(D\mathcal{S}^*_{g_0})$ by (i) and the same construction of $\{g(t)\}$ used in (i) shows that (3.10) holds for any symmetric $(0,2)$ tensor $\hat{h}$ on Ω. Let ψ be any extension of the function $\langle X, \nu_{g_0}\rangle_{g_0}$ to Ω. Let $\hat{h} = \psi G^0$, it then follows from (3.10) that

$$|G^{g_0}|\langle X, \nu_{g_0}\rangle = 0.$$

By Proposition 2.1, we conclude that g_0 is Einstein. □

4. Critical points of $\mathcal{E}_\phi(\cdot)$

Let g_0 be a metric on Ω satisfying **(a1)**. Let H_{g_0} be the mean curvature of Σ in (Ω, g_0). In this section, we discuss the behavior of $\mathcal{E}_\phi(\cdot)$ near g_0 in $\mathcal{M}^\lambda_\gamma$.

Lemma 4.1. *Let $\{g(t)\}_{|t|<\epsilon}$ be a smooth path of metrics with $g(0) = g_0$ and $g(t)|_{T(\Sigma)} = \gamma$ for all t. Let $H(t)$, $A(t)$ denote the mean curvature, the second fundamental form of Σ in $(\Omega, g(t))$, respectively. Let $\mathring{A}(t)$ be the traceless part of $A(t)$. Then*

$$\frac{d}{dt}\mathcal{E}_\phi(g(t))|_{t=0} = \int_\Sigma \phi \left[\frac{n-2}{n-1}H(0)H'(0) - \langle A'(0), \mathring{A}(0)\rangle_\gamma\right] d\gamma_{g_0}.$$

Proof. Since $2H_2 = H^2 - |A|^2$,

$$\begin{aligned}\frac{d}{dt}\mathcal{E}_\phi(g(t))|_{t=0} =& \int_\Sigma \phi\left[H(0)H'(0) - \langle A'(0), A\rangle_\gamma\right] d\gamma_{g_0}\\ =& \int_\Sigma \phi\left[H(0)H'(0) - \frac{H(0)}{n-1}\langle A'(0), \gamma\rangle_\gamma - \langle A'(0), \mathring{A}(0)\rangle_\gamma\right] d\gamma_{g_0}\\ =& \int_\Sigma \phi\left[\frac{n-2}{n-1}H(0)H'(0) - \langle A'(0), \mathring{A}(0)\rangle_\gamma\right] d\gamma_{g_0},\end{aligned}$$

where we also have used the fact $g(t)|_{T(\Sigma)} = \gamma$. □

Lemma 4.2. *Let $\{g(t)\}_{|t|<\epsilon}$ be a smooth path of metrics with $g(0) = g_0$. Let $h = g'(0)$. Let $\{x^1, \ldots, x^{n-1}, x^n\}$ be local coordinates of Ω near Σ such that, when $x^n = 0$, $\{x^1, \ldots, x^{n-1}\}$ form local coordinates on Σ and $\partial_{x^n} = \nu_{g_0}$. Then*

$$A'_{\alpha\beta}(0) = -\frac{1}{2}\left[h_{n\beta;\alpha} + h_{\alpha n;\beta} - h_{\alpha\beta;n}\right] + \frac{1}{2}A_{\alpha\beta}(0)h_{nn}, \tag{4.1}$$

where $1 \le \alpha, \beta \le n-1$.

Proof. The outward unit normal $\nu_{g(t)}$ of Σ in $(\Omega, g(t))$ is given by

$$\nu_{g_t} = \frac{\nabla_{g(t)}x^n}{|\nabla_{g(t)}x^n|_{g(t)}} = \frac{g^{ni}}{\sqrt{g^{nn}}}\partial_{x_i}.$$

Let Γ^k_{ij}, ∇^t denote the Christoffel symbols, the covariant differentiation of $g(t)$, respectively. Then

$$A_{\alpha\beta}(t) = -\langle \nu_{g(t)}, \nabla^t_{\partial_{x^\alpha}}\partial_{x^\beta}\rangle_{g(t)} = -\frac{1}{\sqrt{g^{nn}}}\Gamma^n_{\alpha\beta}.$$

Therefore,

$$\begin{aligned} A'_{\alpha\beta}(0) =& -\frac{1}{2}\frac{1}{\sqrt{g^{nn}}}g^{nj}\left(h_{\alpha j;\beta}+h_{j\beta;\alpha}-h_{\alpha\beta;j}\right)-\frac{1}{2}\frac{g^{nj}g^{in}h_{ij}}{(g^{nn})^{\frac{3}{2}}}\Gamma^n_{\alpha\beta}(0)\\ =& -\frac{1}{2}\left(h_{\alpha n;\beta}+h_{n\beta;\alpha}-h_{\alpha\beta;n}\right)+\frac{1}{2}h_{nn}A_{\alpha\beta}, \end{aligned}$$

which gives (4.1). □

Proposition 4.1. *Suppose ϕ is a nontrivial function on Σ such that*

(i) *$\phi\, H_{g_0}$ is the boundary value of some $N \in Ker(D\mathcal{S}^*_{g_0})$;*

(ii) *Σ is umbilical in (Ω, g_0) at every point $p \in \Sigma$ where $\phi(p) \neq 0$.*

Then $\frac{d}{dt}\mathcal{E}_\phi(g(t))|_{t=0} = 0$ for any path $\{g(t)\} \subset \mathcal{M}^\lambda_\gamma$ with $g(0) = g_0$.

Proof. Condition (ii) simply means $\mathring{A}(0) = 0$ wherever $\phi \neq 0$ along Σ. Hence, by Lemma 4.1,

$$\begin{aligned} \frac{d}{dt}\mathcal{E}_\phi(g(t))|_{t=0} &= \frac{n-2}{n-1}\int_\Sigma \phi H_{g_0}H'(0)d\gamma_{g_0}\\ &= \frac{n-2}{n-1}\frac{d}{dt}\mathcal{E}^{(1)}_{\phi H_{g_0}}(g(t))|_{t=0}, \end{aligned}$$

where the functional $\mathcal{E}^{(1)}_{(\cdot)}(\cdot)$ is defined in Remark 1.1. But

$$\frac{d}{dt}\mathcal{E}^{(1)}_{\phi H_{g_0}}(g(t))|_{t=0} = 0$$

by condition (i) and [10, Th. 2.1]. Hence the result follows. □

Next we want to show that the converse of Proposition 4.1 is also true. This is proved in the next two lemmas.

Lemma 4.3. *If $\frac{d}{dt}\mathcal{E}_\phi(g(t))|_{t=0} = 0$ for any path $\{g(t)\} \subset \mathcal{M}^\lambda_\gamma$ with $g(0) = g_0$, then ϕH_{g_0} is the boundary value of a function in* $\mathrm{Ker}(D\mathcal{S}^*_{g_0})$.

Proof. Let $\hat{h}$ be any symmetric $(0,2)$ tensor with compact support in Ω. By [11, Proposition 1], one can find a positive function $u(x,t) = u_t(x)$, with $x \in \Omega$ and $|t| < \delta$ for some small δ, such that $u_0 = 1$ on Ω, $u_t = 1$ at Σ and $g(t) := u_t^{\frac{4}{n-2}}(g_0 + t\hat{h}) \in \mathcal{M}^\lambda_\gamma$. Let $h = g'(0)$, then $h = \frac{4}{n-2}vg_0 + \hat{h}$, where $v = \frac{\partial u}{\partial t}|_{t=0}$. By Lemma 4.1, this path $\{g(t)\}$ satisfies

$$\int_\Sigma \phi\left[\frac{n-2}{n-1}H_{g_0}H'(0) - \langle A'(0), \mathring{A}(0)\rangle_\gamma\right]d\gamma_{g_0} = 0. \tag{4.2}$$

We claim that, for such a variation, one indeed has

$$\langle A'(0), \mathring{A}(0)\rangle_\gamma = 0. \tag{4.3}$$

If this is true, then (4.2) shows $\frac{d}{dt}\mathcal{E}^{(1)}_{\phi H_{g_0}}(g(t))|_{t=0} = 0$ for this path $\{g(t)\}$. Therefore, by the same proof of [10, Th. 2.1], the result follows.

To prove (4.3), we note that, along Σ,

$$h_{n\beta;\alpha} = \frac{4}{n-2} v_{,\alpha} g_{0 n\beta} + \hat{h}_{n\beta;\alpha} = 0$$

because $\hat{h}$ has compact support and $g_{0 n\beta} = 0$. Here "," denotes partial derivative. Similarly

$$h_{\alpha\beta;n} = \frac{4}{n-2} v_{,n} g_{0\alpha\beta} + \hat{h}_{\alpha\beta;n} = \frac{4}{n-2} v_{,n} \gamma_{\alpha\beta},$$

$$h_{nn} = \frac{4}{n-2} v g_{0 nn} + \hat{h}_{nn} = 0$$

because $v = 0$ at Σ. By Lemma 4.2, we have

$$A'_{\alpha\beta}(0) = \frac{2}{n-2} v_{,n} \gamma_{\alpha\beta}.$$

Thus,

$$\langle A'(0), \mathring{A}(0)\rangle_\gamma = \frac{2}{n-2} v_{,n} \langle \gamma, \mathring{A}(0)\rangle_\gamma = 0$$

as $\mathring{A}(0)$ is traceless. Hence, (4.3) is true. This completes the proof. □

Lemma 4.4. *Suppose $\frac{d}{dt}\mathcal{E}_\phi(g(t))|_{t=0} = 0$ for any path $\{g(t)\} \subset \mathcal{M}^\lambda_\gamma$ with $g(0) = g_0$, Then Σ is umbilical at every point $p \in \Sigma$ with $\phi(p) \neq 0$.*

Proof. By Lemma 4.3, ϕH_{g_0} is the boundary value of a function in $\mathrm{Ker}(D\mathcal{S}^*_{g_0})$. Hence, by [10, Th. 2.1],

$$\int_\Sigma \phi H_{g_0} H'(0) d\gamma_{g_0} = 0$$

for any variation $\{g(t)\} \subset \mathcal{M}^K_\gamma$. Therefore, by Lemma 4.1,

$$\int_\Sigma \phi \langle A'(0), \mathring{A}(0)\rangle_\gamma d\gamma_{g_0} = 0 \tag{4.4}$$

for all such paths.

Now let $p \in \Sigma$ with $\phi(p) > 0$ for instance. Let $\eta \geq 0$ be a smooth function with support near p such that $\eta(p) = 1$ and $\eta(q) > 0$ implies $\phi(q) > 0$. Let $\hat{h}$ be a symmetric $(0,2)$ tensor on Ω such that $\hat{h}|_{T(\Sigma)} = 0$, $\hat{h}(\nu_{g_0}, w) = 0$ for all w tangential to Σ and $\hat{h}(\nu_{g_0}, \nu_{g_0}) = 1$ along Σ. By [11, Proposition 1], there exists a positive function $u(x,t) = u_t(x)$, with $x \in \Omega$ and $|t| < \delta$ for some small δ, such that $u_0 = 1$ on Ω, $u_t = 1$ at Σ and

$$g(t) := u_t^{\frac{4}{n-2}}(g_0 + t\eta\hat{h}) \in \mathcal{M}^\lambda_\gamma.$$

Let $h = g'(0)$, then $h = \frac{4}{n-2} v g_0 + \eta \hat{h}$, where $v = \frac{\partial u}{\partial t}|_{t=0}$. At Σ, we have

$$h_{n\beta;\alpha} = \eta \hat{h}_{n\beta;\alpha} = -\eta \left(\Gamma^k_{\alpha n} \hat{h}_{k\beta} + \Gamma^k_{\alpha\beta} \hat{h}_{nk} \right) = \eta A_{\alpha\beta}(0), \tag{4.5}$$

$$\begin{aligned} h_{\alpha\beta;n} =& \frac{4}{n-2} v_{,n} \gamma_{\alpha\beta} + \eta_{,n} \hat{h}_{\alpha\beta} + \eta \hat{h}_{\alpha\beta;n} \\ =& \frac{4}{n-2} v_{,n} \gamma_{\alpha\beta} - \eta \left(\Gamma^k_{n\alpha} \hat{h}_{k\beta} + \Gamma^k_{\beta n} \hat{h}_{\alpha k} \right) \\ =& \frac{4}{n-2} v_{,n} \gamma_{\alpha\beta}, \end{aligned} \tag{4.6}$$

$$h_{nn} = \frac{4}{n-2} v + \eta = \eta. \tag{4.7}$$

Therefore, by Lemma 4.2 and (4.5) – (4.7),

$$\begin{aligned} A'_{\alpha\beta}(0) =& -\frac{1}{2} \left[2\eta A_{\alpha\beta}(0) - \frac{4}{n-2} v_{,n} \gamma_{\alpha\beta} \right] + \frac{1}{2} \eta A_{\alpha\beta}(0) \\ =& -\frac{1}{2} \left[\eta A_{\alpha\beta}(0) - \frac{4}{n-2} v_{,n} \gamma_{\alpha\beta} \right]. \end{aligned}$$

Plugging it to (4.4) gives

$$\begin{aligned} 0 =& \int_\Sigma \phi \langle \left(\eta A - \frac{4}{n-2} v_{,n} \gamma \right), \mathring{A}(0) \rangle_\gamma d\gamma_{g_0} \\ =& \int_\Sigma \phi \eta \langle \mathring{A}, \mathring{A} \rangle_\gamma d\gamma_{g_0}. \end{aligned}$$

Since $\phi\eta \geq 0$ and $\eta(p)\phi(p) > 0$. We conclude $\mathring{A}(p) = 0$. □

Theorem 1.2 in the introduction now follows directly from Proposition 4.1, Lemma 4.3 and Lemma 4.4.

We end this paper with a proposition relating Theorems 1.1 and 1.2, followed by an example.

Proposition 4.2. *Let (Ω^n, g_0) be a compact Riemannian manifold with boundary Σ. Suppose X is a conformal Killing vector field on (Ω, g_0) such that X does not vanish along Σ and is normal to Σ. Then every point in Σ is umbilical and the mean curvature H_{g_0} of Σ satisfies*

$$H_{g_0} = \frac{n-1}{n} \frac{div_{g_0} X}{\langle X, \nu_{g_0} \rangle}, \tag{4.8}$$

As a result, if $\mathcal{S}(g_0) = \lambda n(n-1)$ and if $\frac{d}{dt} \mathcal{F}_X(g(t))|_{t=0} = 0$ for all variation $\{g(t)\}_{|t|<\epsilon} \subset \mathcal{M}_0^\lambda$, then $\frac{d}{dt} \mathcal{E}_\phi(g(t))|_{t=0} = 0$ for all variation $\{g(t)\}_{|t|<\epsilon} \subset \mathcal{M}_\gamma^\lambda$, where $\phi = \langle X, \nu_0 \rangle$.

Proof. By the assumption on X, $\nu_{g_0} = X/\langle X, \nu_{g_0}\rangle$ at Σ. Let $\{e_i\}_{1\le i\le n}$ be a local orthonormal frame with $e_n = \nu_{g_0}$ at points in Σ. Then the second fundamental form A of Σ is given by

$$A_{\alpha\beta} = \langle \nabla_{e_\alpha}\nu_{g_0}, e_\beta\rangle = \frac{1}{\langle X, \nu_{g_0}\rangle}\langle \nabla_{e_\alpha} X, e_\beta\rangle = \frac{1}{\langle X, \nu_{g_0}\rangle}\eta_{\beta|\alpha}$$

for $1 \le \alpha, \beta \le n-1$, where η is the 1-form dual to X on (Ω, g_0). Hence,

$$A_{\alpha\beta} = \frac{1}{2}(A_{\alpha\beta} + A_{\beta\alpha}) = \frac{1}{2\langle X, \nu_{g_0}\rangle}(\eta_{\alpha|\beta} + \eta_{\beta|\alpha}) = \frac{\mathrm{div}_{g_0} X}{n\langle X, \nu_{g_0}\rangle}\gamma_{\alpha\beta},$$

which proves the first statement of the proposition.

Next, suppose $\mathcal{S}(g_0) = \lambda n(n-1)$ and $\frac{d}{dt}\mathcal{F}_X(g(t))|_{t=0} = 0$ for all variation $\{g(t)\}_{|t|<\epsilon} \subset \mathcal{M}_0^\lambda$. Then, by Theorem 1.1 (i), $\mathrm{div}_{g_0} X \in \mathrm{Ker}(D\mathcal{S}^*_{g_0})$. Hence, by (4.8) and Theorem 1.2, $\frac{d}{dt}\mathcal{E}_\phi(g(t))|_{t=0} = 0$ for all variation $\{g(t)\}_{|t|<\epsilon} \subset \mathcal{M}_\gamma^\lambda$. □

Example Consider a warped product $(\Omega^n, g) = (I \times N, dr^2 + f^2(r)\bar{g})$ where $I \subset \mathbb{R}^1$ is a closed interval, $(N, \bar{g})$ is a closed Riemannian manifold N, and f is a smooth positive function on I. Consider a vector field $X = \phi(r)\partial_r$, where ϕ is any smooth function on I. The 1-form dual to X is $\eta = \phi(r)dr$. Let $\{x^i\}_{1\le i\le n-1}$ be local coordinates of N and let $x^n = r$. The Christoffel symbols $\{\Gamma^n_{ij}\}$ of g are given by

$$\begin{cases} \Gamma^n_{ij} &= -ff'\bar{g}_{ij}, \quad \text{if } 1 \le i, j \le n-1; \\ \Gamma^n_{ni} &= 0, \qquad\quad\ \ \text{if } 1 \le i \le n-1; \\ \Gamma^n_{nn} &= 0. \end{cases}$$

Since $\bar{g}_{ij} = f^{-2}g_{ij}$, we have

$$\eta_{i|j} = \frac{\partial \eta_i}{\partial x^j} - \Gamma^k_{ij}\eta_k = \frac{\partial \eta_i}{\partial x^j} - \Gamma^n_{ij}\phi = \begin{cases} \frac{f'}{f}\phi g_{ij}, & \text{if } 1 \le i, j \le n-1; \\ \phi', & \text{if } i = j = n; \\ 0, & \text{otherwise.} \end{cases}$$

From this, one concludes that $\eta_{i|j} + \eta_{i|j} = \frac{2}{n}(\mathrm{div}_g X)g_{ij}$, $1 \le i, j \le n$, if and only if $\phi' = (f'\phi)/f$, which is equivalent to $\phi = cf$ for some constant c. Now set $\phi = f$, then $\mathrm{div}_g X = nf'$. Note that this X is never zero on $\partial\Omega$ and is normal to $\partial\Omega$. Classification for f' being in $\mathrm{Ker}(D\mathcal{S}*_g)$ can be found in [8, 9].

References

[1] Arnowitt, R.; Deser, S., and Misner, C. W., *Coordinate invariance and energy expressions in general relativity*, Phys. Rev. (2) **122** (1961), 997–1006.

[2] Ashtekar, A.; Hansen, R. O., *A unified treatment of null and spatial infinity in general relativity. I. Universal structure, asymptotic symmetries, and conserved quantities at spatial infinity*, J. Math. Phys. **19** (1978), no. 7, 1542–1566.

[3] Corvino, J., *Scalar curvature deformation and a gluing construction for the Einstein constraint equations*, Comm. Math. Phys. **214** (2000), 137–189.
[4] Chruściel, P., *A remark on the positive-energy theorem*, Class. Quantum Grav. **3** (1986), L115–L121.
[5] Chruściel, P.; Herzlich, M., *The mass of asymptotically hyperbolic Riemannian manifolds*, Pacific J. Math. **212** (2003), no. 2, 231–264.
[6] Herzlich, M., *Computing asymptotic invariants with the Ricci tensor on asymptotically flat and hyperbolic manifolds*, arXiv:1503.00508.
[7] Huang, L.-H., *On the center of mass in General Relativity*, in Fifth International Congress of Chinese mathematicians, Part I, 2, AMS/IP Stud. Adv. Math., vol. 51, Amer. Math. Soc., Providence, RI, 2012, 575–591.
[8] Kobayashi, O., *A differential equation arising from scalar curvature function*, J. Math. Soc. Japan 34 (1982), no. 4, 665–675.
[9] Kobayashi, O.; Obata, M., *Conformally-flatness and static space-time*, Manifolds and Lie groups (Notre Dame, Ind., 1980), pp. 197–206, Progr. Math., **14**, Birkhäuser, Boston, Mass., 1981.
[10] Miao, P.; Shi, Y.-G, and Tam, L.-F., *On geometric problems related to Brown-York and Liu-Yau quasilocal mass*, Comm. Math. Phys. **298** (2010), no. 2, 437–459.
[11] Miao, P.; Tam, L.-F., *On the volume functional of compact manifolds with boundary with constant scalar curvature*, Calc. Var., (2009) **36**:141–171.
[12] Miao, P.; Tam, L.-F., *Evaluation of the ADM mass and center of mass via the Ricci tensor*, Proc. Amer. Math. Soc. **144** (2016), no 2, 753–761.
[13] Miao, P.; Tam; L.-F., and Xie, N.-Q., *Quasi-local mass integrals and the total mass*, arXiv:1510.07756.
[14] Wang, G.; Wu, J., *Chern's magic form and the Gauss-Bonnet-Chern mass*, arXiv:1510.03036.

(Pengzi Miao) Department of Mathematics, University of Miami, Coral Gables, FL 33146, USA.
E-mail address: pengzim@math.miami.edu

(Luen-Fai Tam) The Institute of Mathematical Sciences and Department of Mathematics, The Chinese University of Hong Kong, Shatin, Hong Kong, China.
E-mail address: lftam@math.cuhk.edu.hk